maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comment arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 2007	2 DEDORT TYPE			3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
The STEREO Mission: A Three-Dimensional View of the Sun and Heliosphere				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,4555 Overlook Avenue SW,Washington,DC,20375				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ	ABILITY STATEMENT ic release; distributi	on unlimited			
13. SUPPLEMENTARY NO	OTES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	3	

Report Documentation Page

Form Approved OMB No. 0704-0188

The STEREO Mission: A Three-Dimensional View of the Sun and Heliosphere

J.W. Cook, J.S. Newmark, and R.A. Howard Space Science Division

Coronal Mass Ejections: Coronal mass ejections (CMEs) are massive outflows of material from the Sun's corona, traveling into the surrounding heliosphere at speeds up to 1000–2000 kilometers per second. When directed toward the Earth, CMEs can result in damage to satellites, enhanced auroral displays, blackouts of power grids on Earth, and interference with surface radio communications. The first CME detected from space was observed by an NRL coronagraph onboard the Seventh Orbiting Solar Observatory (OSO-7) in December 1971.¹

Much of our recent knowledge of CMEs has been obtained from an NRL visible-light coronagraph (LASCO, the Large Angle Spectroscopic Coronagraph²) on the Solar and Heliospheric Observatory (SOHO) spacecraft, launched in 1995. But SOHO observes from only one viewpoint, leaving three-dimensional structure ambiguous for individual CMEs. The next step beyond SOHO is the Solar Terrestrial Relations Observatory (STEREO) mission, consisting of two spacecraft observing the Sun from two different viewpoints, allowing a three-dimensional view of the initiation of CMEs and their propagation outwards. In addition, STEREO will carry a new type of heliospheric coronagraph that is off-pointed from the solar disk, and can observe from the side the Sun-Earth line all the way out to Earth. The STEREO mission is meant to increase our understanding of such topics as solar structures and their properties involved in CME initiation; threedimensional structure and kinematics of CMEs; threedimensional structure of active regions, coronal loops, and streamers; propagation of CMEs into the corona and interplanetary medium; and the effects of CMEs through the heliosphere to the Earth.

SECCHI Instruments and Their Development:

The Sun Earth Connection Coronal and Heliospheric Investigation (SECCHI) experiment³ is one component of STEREO. SECCHI was built by an international consortium led by the Naval Research Laboratory. Major components were developed at NRL, Lockheed Martin Solar and Astrophysics Laboratory (Palo Alto, CA), Goddard Space Flight Center (Greenbelt, MD), University of Birmingham (UK), Rutherford Appleton Laboratory (UK), Max-Planck-Institut für Sonnensystemforschung (Germany), Centre Spatiale de Liège (CSL, Belgium), Institut d'Optique (France), and Institut d'Astrophysique Spatiale (France).

Each of the twin SECCHI instrument suites consists of five telescopes. Figure 4 illustrates how each telescope contributes to the total coverage (using existing data from SOHO observations to mimic the SECCHI observations). An extreme-ultraviolet imager (EUVI) will image the solar transition region and low corona, in four emission lines, on the disk and out to 1.7 R₀ (solar radii). The COR1 and COR2 telescopes, visible light Sun-centered coronagraphs with nested fields of view, image the outer corona over 1.4–15 R₀. They were split into two telescopes for better coverage of the decreasing coronal brightness over this range. The Heliospheric Imager (HI) coronagraph is offpointed from the solar disk, and can observe from the side a distance range from 15 R₀ out to the radius of Earth at 215 R₀. Again, coverage is divided between two telescopes, HI-1 and HI-2. A series of linear occulters shelter HI from the glare of the Sun, while internal concentric curved baffles suppress stray light. A Guide Telescope (GT) serves as the spacecraft fine Sun sensor and provides the error signal for the EUVI fine pointing system. The SECCHI Electronics Box (SEB) was developed by the Space Electronics Systems Development Branch at NRL. Figure 5 illustrates the SECCHI instrument suite in the laboratory.

SECCHI went through an instrument development process that included initial design, multiple project reviews, fabrication, integration to two complete packages, and laboratory testing. Characterization of the COR1, COR2, EUVI, and GT was performed at NRL, measuring stray light, spatial resolution, polarization, and photometric response. HI was calibrated at CSL, Belgium. Instrument integration and environmental testing (thermal vacuum, vibration, and electromagnetic interference and compatibility) were performed at NRL using Naval Center for Space Technology facilities prior to spacecraft integration.

Launch and Early Operations: The twin NASA STEREO spacecraft (A and B) were launched on a single Delta rocket from Cape Canaveral on October 25, 2006. Using a lunar encounter, each spacecraft was inserted into an individual final orbit, one spacecraft (B) gradually falling behind Earth in orbit and the other (A) increasingly ahead of Earth. SECCHI first light for each of the A and B telescopes was achieved during the first half of December. Figure 6 illustrates how the scientific observations will be undertaken, showing a simulated Earth-directed CME observed from the A and B spacecraft. The SECCHI COR2A coronagraph already observed its first CME during earliest instrument open-door operations in December 2006.

[Sponsored by NASA and ONR]

References

- ¹R. Tousey, *Space Research XIII*, 713 (1973); and see frontispiece to *Solar Physics* **24**(2) (1972).
- ² G.E. Brueckner, R.A. Howard, M.J. Koomen, C.M. Korendyke, D.J. Michels, J.D. Moses, D.G. Socker, K.P. Dere, P.L. Lamy, A. Llebaria, M.V. Bout, R. Schwenn, G.M. Simnett, D.K. Bedford, and C.J. Eyles, "The Large Angle Spectroscopic Coronagraph (LASCO)," Solar Physics 162, 357 (1995).
- ³ R.A. Howard, J.D. Moses, A. Vourlidas, J.S. Newmark, D.S. Socker, S.P. Plunkett, C.M. Korendyke, J.W. Cook, A. Hurley, J.M. Davila, W.T. Thompson, O.C. St Cyr, E. Mentzell, K. Mehalick, J.R. Lemen, J.P. Wuelser, D.W. Duncan, T.D. Tarbell, R.A. Harrison, N.R. Waltham, J. Lang, C.J. Davis, C.J. Eyles, J.P. Halain, J.M. Defise, E. Mazy, P. Rochus, R. Mercier, M.F. Ravet, F. Delmotte, V. Bothmer, D. Wang, N. Rich, S. Cooper, V. Stephens, G. Maahs, R. Baugh, and D. McMullin, "Sun Earth Connection Coronal and Heliospheric Investigation (SECCHI)," *Space Science Reviews*, in press. ⋄

SECCHI on STEREO: Exploration of the Corona, CMEs, and the Heliosphere From Multiple Viewpoints **SECCHI Science Goals** e IX. X He II · Study structures and their properties involved in CME initiation · Study the three-dimensional structure and kinematics of CMEs Study the three-dimensional structure of active regions, coronal loops, and streamers Study the propagation of CMEs into the corona and interplanetary medium Fe XV Study the effects of CMEs through the heliosphere to the Earth EUVI emission line intensity of COR1 white light polarized structures in the transition region intensity of the corona and inner corona (FOV<1.7 R_®) (FOV 1.4-4.0 R_®) HI2 COR2 38.4 Re 138.7 R. COR1 52 4 R. 200 R_s HI-2 white light total intensity of the HI-1 white light total intensity of the COR2 white light polarized intensity heliosphere (FOV 138.7 R_® centered at heliosphere (FOV 38.4 R_® centered of the corona (FOV 2.5-15.5 R_m) 200 R_®) at 52.4 Re)

FIGURE 4SECCHI telescopes and their fields of view.



FIGURE 5 SECCHI instrument suite in the laboratory.




FIGURE 6The STEREO A and B spacecraft observing an Earth-directed CME.